

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) Nov 2014		2. REPORT TYPE Viewgraph		3. DATES COVERED (From - To) Nov 2014- Dec 2014	
4. TITLE AND SUBTITLE Advanced Multifluid and Collisional-Radiative models for Laser-Plasma Interaction				5a. CONTRACT NUMBER In-house	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Hai Le, Robert Martin, David Bilyeu, Jean-Luc Cambier				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER Q02Z	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRS 1 Ara Drive. Edwards AFB CA 93524-7013				8. PERFORMING ORGANIZATION REPORT NO.	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2014-363	
12. DISTRIBUTION / AVAILABILITY STATEMENT Distribution A: Approved for Public Release; Distribution Unlimited.					
13. SUPPLEMENTARY NOTES Briefing Charts presented at Plasma and Electroenergetics Program Review, Arlington, CA, 18-19 December 2014. PA#14613					
14. ABSTRACT N/A					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 23	19a. NAME OF RESPONSIBLE PERSON Jean-Luc Cambier
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NO (include area code) 661-275-5649



Advanced Multifluid and Collisional-Radiative models for Laser-Plasma Interaction

*AFOSR Plasma and Electroenergetics Review Meeting
December 2014*



Collaborators:

- R. Caflisch (UCLA)
- S. Gimelshein (USC)
- A. Karagozian (UCLA)

Hai Le¹, Robert Martin¹, David Bilyeu²

Jean-Luc Cambier³

¹ERC Inc.

²NRC

³AFRL/RQRS

Distribution A – Approved for public release; Distribution Unlimited. PA# XXXXX



TOC



- **Motivation**
- **Plasma M&S & CR kinetics**
- **Level grouping**
- **Multi-fluid equations & LPI**
- **Beyond translational equilibrium**
- **Summary & future work**



Spacecraft Plasma M&S




- Applications: Hall thrusters, FRC, LIBS, Laser Plasma Interaction, Plasma Discharges
- Complex physics: excitation/ionization, transport, radiation, material, etc.
- Multiple spatial-temporal and density scales.



Current focus:

Develop advanced multiscale algorithms for plasma M&S in highly non-equilibrium condition and with collisional-radiative kinetics


Laser Plasma Interaction




EP Plumes



Ionizing shocks



FRC



Chamber Environment



Plasma M&S



- **Kinetic equation:**

$$\partial_t f + \vec{v} \cdot \nabla_{\vec{x}} f = \frac{1}{\varepsilon} Q(f, f) + Q^{CR}(f, f)$$

Boltzmann Coll. Op.:
$$Q(f, f) = \int_{R^3} \int_{S^2} \sigma(|v - v_*|, \omega) [f(v') f(v_*') - f(v) f(v_*)] d\omega dv'$$

Equilibrium vdf:
$$Q(f, f) = 0 \rightarrow f(v) = \frac{\rho}{(2\pi T)^{3/2}} \exp\left(-\frac{|v - u|^2}{2T}\right)$$

- Fluid regime: $\varepsilon \ll 1$
- Kinetic regime: $\varepsilon = O(1)$
- Collisional plasma: excitation/ionization, CE collisions, radiation, etc.

- **Methods: moment method, PIC, DNS.**

- **Challenges:**

- Multiple species: $f \rightarrow f_s \quad Q(f, f) \rightarrow \sum_t Q(f_s, f_t)$
- Dynamical regime: $\varepsilon \rightarrow \varepsilon(p, x, t)$




Collisional-Radiative (CR) model



- **Non-equilibrium modeling of the atomic state distribution function (ASDF)**
 - Detailed state-to-state model of atomic transition, i.e., excitation, ionization, line radiation, etc.
 - Rates derived based on ab initio cross section.
- **Examples: hypersonic shocks in Ar & N₂**
- **Complications:**
 - Accuracy can require many states
 - Translational nonequilibrium

Atomic CR
Ar shock (31 levs)





Maxwellian CR

- **Hydrogen model:** $E_n = I_H (1 - 1/n^2)$ $I_n = I_H / n^2$ $g_n = n^2$

- **Analytical rates:**

— Excitation/deexcitation:

$$\alpha_{(m|n)}^e \simeq \left[4\pi a_0^2 \cdot \frac{32}{\pi\sqrt{3}} \cdot \bar{v}_e \right] \frac{e^{-x_{nm}}}{n^5 m^3 (n^{-2} - m^{-2})^5},$$

$$\beta_{(n|m)}^e \simeq \left[4\pi a_0^2 \cdot \frac{32}{\pi\sqrt{3}} \cdot \bar{v}_e \right] \frac{1}{n^3 m^5 (n^{-2} - m^{-2})^5}.$$

— Ionization/recombination: $\alpha_{(+|n)}^e \simeq (4\pi a_0^2) \left(\frac{8kT_e}{\pi m_e} \right)^{1/2} n^4 e^{-x_n}$

$$\beta_{(n|+)}^e \simeq \left[\frac{4}{\pi} \frac{a_0^2 h^3}{m_e^2 k T_e} \right] n^6$$



— Line radiation

$$A_{(n|m)} = \left(\frac{8\pi^2 e^2}{m_e c^3} \right) \frac{g_n}{g_m} f_{nm} = \frac{1.6 \times 10^{10}}{m^3 n (m^2 - n^2)} \text{ s}^{-1}$$

- **Rate equation:**

$$\begin{aligned} \frac{dN_n}{dt} = & - \sum_{m>n} \alpha_{(m|n)} N_e N_n + \sum_{m>n} \beta_{(n|m)} N_e N_m + \sum_{m>n} A_{(n|m)} N_m \\ & + \sum_{m<n} \alpha_{(n|m)} N_e N_m - \sum_{m<n} \beta_{(m|n)} N_e N_n - \sum_{m<n} A_{(m|n)} N_n \\ & - \alpha_{(+|n)} N_e N_n + \beta_{(n|+)} N_+ N_e^2. \end{aligned} \quad (8)$$

$$\frac{dN_+}{dt} = \sum_n \alpha_{(+|n)} N_e N_n - \sum_n \beta_{(n|+)} N_+ N_e^2.$$





Level grouping

- **CR modeling: level-grouping** $\rightarrow \mathcal{N}_n = N_{n0} \sum_{i \in n} \frac{N_i}{N_{n0}} \simeq \frac{N_{n0}}{g_{n0}} \underbrace{\sum_{i \in n} g_i e^{-\Delta E_i / T_n}}_{Z_n}$
 - Group effective rates of change
$$\frac{d\mathcal{N}_n}{dt} = -N_e \mathcal{N}_n \left[\sum_{m>n} \sum_{i \in n} \frac{g_i e^{-\Delta E_i / T_n}}{Z_n} \sum_{j \in m} \alpha_{(j|i)} + \sum_{m<n} \sum_{i \in n} \frac{g_i e^{-\Delta E_i / T_n}}{Z_n} \sum_{j \in m} \beta_{(j|i)} \right]$$
 - Internal structure of group is assumed Boltzmann (T_n)
 - Piecewise exponential
 - This does NOT mean the entire ASDF is Boltzmann!!
 - Group temperature must be determined \rightarrow additional conservation equation, e.g.:
$$\frac{d\mathcal{E}_n}{dt} = -N_e \mathcal{N}_n \left[\sum_{m>n} \sum_{i \in n} \frac{g_i e^{-\Delta E_i / T_n}}{Z_n} \sum_{j \in m} E_i \alpha_{(j|i)} + \sum_{m<n} \sum_{i \in n} \frac{g_i e^{-\Delta E_i / T_n}}{Z_n} \sum_{j \in m} E_i \beta_{(j|i)} \right]$$
 - Procedure?

– Solve:

$$\langle \Delta E \rangle_n(T_n^*) + C_v(T_n^*) \delta T_n^* = \frac{\Delta \mathcal{E}_n^{(k)}}{\mathcal{N}_n^{(k)}}$$

with:

$$\Delta \mathcal{E}_n = \sum_{i \in n} (E_i - E_{n0}) N_i = \mathcal{N}_n \langle \Delta E \rangle_n$$

$$C_v(T_n) = \frac{d}{dT_n} \langle \Delta E \rangle_n$$

Flowchart labels: iterated, computed, tabulated

– **However...** $\langle \Delta E \rangle_n \simeq o(\epsilon)$ where $\epsilon = e^{-\Delta E_1 / T_n} \rightarrow \delta T_n^* = o(\epsilon) / o(\epsilon)$

$C_v(T_n) \simeq o(\epsilon)$



Level grouping

- **CR modeling: level-grouping**

- Other approaches?

- Sub-partitioning: lowest level n_0 and total \mathcal{N}_n (no need for \mathcal{E}_n)

$$\delta T_n^* \simeq \frac{T_n^{*2}}{\mathcal{Z}_n(T_n^*) \langle \Delta E \rangle_n(T_n^*)} \left[\frac{\mathcal{N}_n^{(k)}}{N_{n_0}^{(k)}} g_{n_0} - \mathcal{Z}_n(T_n^*) \right] = o(\epsilon)/o(\epsilon) \text{ ...fails}$$

- Sub-partitioning: lowest level n_0 and upper distribution \mathcal{N}_n ,

$$\delta T_n^* \simeq \frac{T_n^{*2}}{\mathcal{Z}'_n(T_n^*) \langle \Delta E \rangle_n(T_n^*)} \left[\frac{\mathcal{N}_n^{(k)}}{N_{n_0}^{(k)}} g_{n_0} - \mathcal{Z}'_n(T_n^*) \right] = o(\epsilon)/o(\epsilon) \text{ ...fails}$$

- Approximate \mathcal{Z}_n by expanding around mean energy: $\overline{\Delta E}_n = \frac{1}{g_n} \sum_{i \in n} g_i \Delta E_i$.

$$\mathcal{Z}_n(T_n) = e^{-\overline{\Delta E}_n/T_n} \sum_{i \in n} g_i \left[1 - \cancel{\frac{\delta_i}{T_n}} + \frac{1}{2} \frac{\delta_i^2}{T_n^2} + \dots \right] \quad \text{where } \delta_i \equiv \Delta E_i - \overline{\Delta E}_n$$




- With n_0 , \mathcal{N}_n partitioning: $1/\ln(1+\epsilon)$...fails
- With n_0 , \mathcal{N}_n partitioning: $1/\ln(\epsilon)$...succeeds!
- Improve with successive iterations... $\mathcal{Z}'_n(T_n) = \bar{g}'_n(T_n) e^{-\overline{\Delta E}'_n/T_n}$



Level grouping: Numerical test



- Isothermal heat bath:**



Works very well (also tested in cooling regime).
Much better than uniform (standard) grouping.



Level grouping: Energy conservation



- **Energy conservation:**

- Conservation follows from definition $\mathcal{E}_n = \bar{E}_n \mathcal{N}_n$

- Start with:

$$\frac{d\Delta\mathcal{E}_n}{dt} \equiv \sum_{i \in n} \Delta E_i \frac{dN_i}{dt} = \frac{d}{dt} (\mathcal{N}'_n \langle \Delta E \rangle_{n'}) = \langle \Delta E \rangle_{n'} \frac{d\mathcal{N}'_n}{dt} + \mathcal{N}'_n \frac{d\langle \Delta E \rangle_{n'}}{dt}$$

$\Delta(\text{overall group population}) : \mathcal{N}_n$
 $\Delta(\text{internal structure}) := C_{v,n'} \frac{dT_n}{dt}$

- Express in terms of conserved variables:

$$\frac{d\mathcal{E}_n}{dt} = [E_{n_0} - \omega_{n'}] \frac{dN_{n_0}}{dt} + [E_{n_0} + \langle \Delta E \rangle_{n'} + \xi_{n'}] \frac{d\mathcal{N}'_n}{dt} \quad \text{with} \quad \xi_{n'} = \frac{C_{v,n'} T_n^2}{\left(\langle \Delta E \rangle_{n'} + T_n^2 \frac{d \ln \bar{z}_n}{dT_n} \right)} \quad \text{and} \quad \omega_{n'} = \xi_{n'} \frac{\mathcal{N}_{n'}}{N_{n_0}}$$



Level grouping: Energy conservation



- **Energy conservation:**

- Finally...Procedure shown to be equivalent to replacing energies by “effective” (condition-dependent) values (\approx EOS)

Excitation: $\bar{\alpha}_{(m_0|n_0)}^E = (\bar{E}_{m_0} - \bar{E}_{n_0}) \cdot \bar{\alpha}_{(m_0|n_0)}$

$$\bar{\alpha}_{(m'|n_0)}^E = (\bar{E}_{m'} - \bar{E}_{n_0}) \cdot \bar{\alpha}_{(m'|n_0)}$$

$$\bar{\alpha}_{(m_0|n')}^E = (\bar{E}_{m_0} - \bar{E}_{n'}) \cdot \bar{\alpha}_{(m_0|n')}$$

$$\bar{\alpha}_{(m'|n')}^E = (\bar{E}_{m'} - \bar{E}_{n'}) \cdot \bar{\alpha}_{(m'|n')}$$

Ionization: $\bar{\alpha}_{(+|n_0)}^E = (I_H - \bar{E}_{n_0}) \cdot \bar{\alpha}_{(+|n_0)}$

$$\bar{\alpha}_{(+|n')}^E = (I_H - \bar{E}_{n'}) \cdot \bar{\alpha}_{(+|n')}$$


with

$$\bar{E}_{n_0} = E_{n_0} - \omega_{n'}$$

$$\bar{E}_{n'} = E_{n_0} + \langle \Delta E \rangle_{n'} + \xi_{n'}$$

- **NOW**, energy is conserved

(down to round-off) →





Multi-fluid equations

- **Multi-fluid model:**

- 5-moment:

$$\partial_t \rho_s + \nabla \cdot (\rho_s \mathbf{u}_s) = \omega_s^\rho$$

$$\partial_t (\rho_s \mathbf{u}_s) + \nabla \cdot (\rho_s \mathbf{u}_s \mathbf{u}_s + \mathbb{P}_s) = Z_s e n_s (\mathbf{E} + \mathbf{u}_s \times \mathbf{B}) + \mathbf{f}_s^m$$

$$\partial_t \epsilon_s + \nabla \cdot (\mathbf{u}_s \epsilon_s + \mathbb{P}_s \cdot \mathbf{u}_s) + \nabla \cdot \mathbf{q}_s = \mathbf{j}_s \cdot \mathbf{E} + \omega_s^\epsilon$$

- Add Maxwell's equations:

$$\begin{aligned} \nabla \times \mathbf{E} &= -\partial_t \mathbf{B} & \nabla \cdot \mathbf{E} &= \frac{e}{\epsilon_0} (Z_i n_i - n_e) \\ \nabla \times \mathbf{B} &= \mu_0 \mathbf{j} - \frac{1}{c^2} \partial_t \mathbf{E} & \nabla \cdot \mathbf{B} &= 0 \end{aligned}$$

- Add collisions:

- Elastic – Braginskii terms
 - Inelastic – Rates depend on both T and relative velocity


$$k_i = n_n n_e \int \int f_n f_e g \sigma_i''(g; \Omega_1, \Omega_2) d\Omega_1 d\Omega_2 d^3 v_n d^3 v_e \quad \Rightarrow \quad k_i = k_i(T_e, |\mathbf{u}_n - \mathbf{u}_e|)$$

- Multi-fluid CR model from fundamental principles being developed (incl. detailed balance)



Multi-fluid equations

- Electromagnetic shock: generalized Brio-Wu¹
 - FV with WENO reconstruction and RK3



¹Shumlak & Loverich, JCP 2003



Two-Fluid model

- Assume fully ionized plasma, electrostatic field:

$$\partial_t \rho_e + \nabla \cdot (\rho_e \mathbf{u}_e) = 0$$

$$\partial_t \rho_i + \nabla \cdot (\rho_i \mathbf{u}_i) = 0$$

$$\partial_t (\rho_e \mathbf{u}_e) + \nabla \cdot (\rho_e \mathbf{u}_e \mathbf{u}_e + p_e \mathbb{I}) = -en_e \mathbf{E} + \rho_e \nu_{ei} \mathbf{W}_{ei} + \mathbf{f}_p$$

$$\partial_t (\rho_i \mathbf{u}_i) + \nabla \cdot (\rho_i \mathbf{u}_i \mathbf{u}_i + p_i \mathbb{I}) = Z_i en_i \mathbf{E} - \rho_e \nu_{ei} \mathbf{W}_{ei}$$

$$\partial_t E_e + \nabla \cdot [(E_e + p_e) \mathbf{u}_e] = -\nabla \cdot \mathbf{q}_e + \mathbf{j}_e \cdot \mathbf{E} + \rho_e \nu_{ei} \mathbf{W}_{ei} \cdot \bar{\mathbf{u}}_{ei}$$

$$+ 3n_e \nu_{ei} k(T_i - T_e) + \mathbf{f}_p \cdot \mathbf{u}_e + W_L$$

$$\partial_t E_i + \nabla \cdot [(E_i + p_i) \mathbf{u}_i] = \mathbf{j}_i \cdot \mathbf{E} - \rho_e \nu_{ei} \mathbf{W}_{ei} \cdot \bar{\mathbf{u}}_{ei} - 3n_e \nu_{ei} k(T_i - T_e)$$

- Gauss's law for electrostatic field $\nabla \cdot \mathbf{E} = \frac{e}{\epsilon_0} (Z_i n_i - n_e)$

- Transport: Spitzer with flux limiter

$$\mathbf{q}_e = \min\left(f \frac{q_{FS}}{|\mathbf{q}_{SH}|}, \mathbf{q}_{SH}\right)$$

$$\kappa_e = \frac{\gamma_Z n_e k^2 T_e}{m_e \nu_{ei}}$$

$$\mathbf{q}_{SH} = -\kappa_e \nabla T_e$$

$$\gamma_Z \approx \frac{3.22554(Z_i + 0.24)}{1 + 0.24 Z_i}$$

$$q_{FS} = \left(\frac{2}{\pi}\right)^{1/2} k T_e v_{Te}$$



Electrodynamics

- Maxwell's equations for laser E and B fields (different from plasma)

$$\partial_{tt}\mathbf{E} - c^2\nabla^2\mathbf{E} + c^2\nabla(\nabla \cdot \mathbf{E}) + \frac{1}{\epsilon_0}\partial_t\mathbf{j} = 0$$
$$\partial_{tt}\mathbf{B} - c^2\nabla^2\mathbf{B} - \frac{1}{\epsilon_0}\nabla \times \mathbf{j} = 0$$

- Cold plasma response via Ohm's law:

$$\partial_t\mathbf{j} + \nu\mathbf{j} = \epsilon_0\omega_p^2\mathbf{E}$$

- Assuming monochromatic wave:

$$\mathbf{E}(\mathbf{x}, t) = \hat{\mathbf{E}}(\mathbf{x})e^{-i\omega t} + c.c.$$

$$\mathbf{B}(\mathbf{x}, t) = \hat{\mathbf{B}}(\mathbf{x})e^{-i\omega t} + c.c.$$

$$\mathbf{j}(\mathbf{x}, t) = \hat{\mathbf{j}}(\mathbf{x})e^{-i\omega t} + c.c.$$

$$\hat{\mathbf{j}} = \sigma\hat{\mathbf{E}}$$

$$\sigma = \frac{i\epsilon_0\omega_p^2}{\omega(1 + i\nu/\omega)}$$

Complex conductivity

Dielectric function

$$\epsilon = \eta^2 = 1 - \frac{\omega_p^2}{\omega^2(1 + i\nu/\omega)}$$

Refractive index

- Wave equations becomes:

$$\nabla^2\hat{\mathbf{E}} + k_0^2\epsilon\hat{\mathbf{E}} - \nabla(\nabla \cdot \hat{\mathbf{E}}) = 0$$

$$\nabla^2\hat{\mathbf{B}} + k_0^2\epsilon\hat{\mathbf{B}} + \nabla(\ln \epsilon) \times (\nabla \times \hat{\mathbf{B}}) = 0$$



Ponderomotive force and collisional heating



- **Nonlinear force (Hora, *PoF* 1985)**

- General: $\mathbf{f}_{nl} = \mathbf{j} \times \mathbf{B} + \epsilon_0 \mathbf{E} \nabla \cdot \mathbf{E} + \epsilon_0 (1 + \omega^{-1} \partial_t) \nabla \cdot (\eta^2 - 1) \mathbf{E} \mathbf{E}$
- Assume steady-state and time-average: ponderomotive force

$$\mathbf{f}_p = \langle \mathbf{f}_{nl} \rangle = -\nabla \cdot [\langle \mathbf{T} \rangle - \epsilon_0 (\eta^2 - 1) \langle \mathbf{E} \mathbf{E} \rangle]$$

↑
Maxwell tensor

- **Collisional absorption:**


- Poynting theorem: $\nabla \cdot \mathbf{S} = -\mathbf{j} \cdot \mathbf{E}$

- Time-average: $W_L = \langle \mathbf{j} \cdot \mathbf{E} \rangle = -\nabla \cdot \langle \mathbf{S} \rangle$

↑
Poynting vector

- **Assume 1-d propagation:**


$$\mathbf{f}_p = -\frac{\epsilon_0 \omega_p^2}{4\omega^2} \partial_x (\hat{\mathbf{E}} \hat{\mathbf{E}}^*)$$





Laser-Plasma Interaction

- Ion acceleration due to ponderomotive force






Beyond Maxwellian: $Kn = O(1)$



- **Not-too-far from equilibrium (isotropic)**
 - Discretized EEDF yields rate equations for discrete elements (“bin”)
 - DB enforced at microscopic level
 - High-order, implicit and energy conserving
 - More efficient compared to MCC.
- **Far from equilibrium**
 - MCC algorithm for inelastic collisions
 - Can resolve anisotropic vdf
 - Drawback: slow convergence, reaction branching, singular rates, computational particle growth



$$\bar{n}_i = N_e \int_{\varepsilon_i}^{\varepsilon_i + \Delta\varepsilon} d\varepsilon f_e(\varepsilon)$$

$$\frac{d\bar{n}_i}{dt} = -N_l \bar{n}_i \sum_j \bar{k}_{(j|i)}^{\text{exc}} + N_u \sum_j \bar{n}_j \bar{k}_{(i|j)}^{\text{dex}}$$



$$\frac{d\bar{n}_j}{dt} = +N_l \sum_i \bar{n}_i \bar{k}_{(j|i)}^{\text{exc}} - N_u \bar{n}_j \sum_i \bar{k}_{(i|j)}^{\text{dex}}$$



Beyond Maxwellian: Particle Merging




- Scheme built on merging 3+ particles to 2.
- Mass, momentum and kinetic energy are exactly conserved; Electrostatic energy also conserved in physical space.
- Split analogously defined by merging only fractions of original particles.
- Octree in velocity space inhibits thermalization by ensuring only near neighbor particles are merged.
- Higher-moment conserving schemes have been obtained with increased number of merge result particles generated.





Particle Merge: Importance of Octree






Particle Merge: DC Breakdown Case



- Tested in 3D ES-PIC of 1-KV DC Breakdown
- MCC-Ionization, Chain Branching & Cathode Secondary Emission causes exponential growth in cost
- With Merging: Density matches despite vastly different number of Computational Particles/Cell
- Negligible overhead demonstrated in comparison of wall-clock/iteration with merge every iteration
- Enables direct control of computational cost in particle methods
- Future Work: Test merge in non-Maxwellian laser plasma test case





Conclusion



- **Multiscale algorithms for nonequilibrium flows with CR kinetics**
 - Level grouping schemes of electronic states of atoms.
 - Multi-fluid equations developed to efficiently capture electron “hydrodynamics”
 - Particle merge/split for particle management, efficient sampling, inelastic collisions ...
- **Ongoing works:**
 - Multi-D simulation with level grouping
 - Modeling of inelastic collisions in multi-fluid
 - High-order particle merging schemes